

PRE-SURGERY PREPARATION

DO NOT take aspirin or any anti-inflammatory medications such as ibuprofen, Motrin, Aleve or Advil for 2 weeks prior to and 1 week after your surgery.

STOP taking vitamin E and fish oil supplements for 2 weeks prior to and 2 weeks after surgery. A multivitamin is fine as long as it has less than 700 IU per day.

STOP smoking for at least 2 weeks before and 2 weeks after your surgery. Smoking greatly increases your chance of complications.

MAKE ARRANGEMENTS to have someone drive you to and from surgery and to your office visit the day after surgery.

ARRANGE for someone to care for you the first 24-72 hours after surgery.

FILL your prescriptions before surgery. It is not necessary to bring them with you to the surgery center.

SET UP a home recovery area with lots of pillows in a quiet area where you can rest. A recliner is very helpful if you have one.

HAVE light foods on hand such as saltine crackers, protein shakes, soup, applesauce, jell-o, popsicles, etc. Also have plenty of fluids available. Water, dilute fruit juices, dilute Gatorade and caffeine free tea are good choices.

DO NOT eat or drink anything after 12:00 midnight. This includes water, candy and chewing gum. We suggest having a light dinner no later than 7 pm the evening before your surgery to minimize the chance of nausea. Drinking plenty of water (64oz/day) or more for two days prior to your surgery will also help prevent post-op nausea.

WEAR only comfortable, loose-fitting clothes that open in the front to the surgery center. Slippers or slip-on shoes are preferred. For patients having a tummy tuck a robe is best.

DO NOT wear makeup, jewelry, contact lenses or body piercings to the surgery center.

SHOWER and wash your hair the morning of surgery. Wash the area you are having surgery on with Hibiclens (antiseptic cleanser), which will be provided for you at your pre-op appointment. Do not apply any powder, lotion or deodorant after you shower.



HEDIEH A. STEFANACCI, M.D.
AESTHETIC & RECONSTRUCTIVE PLASTIC SURGERY

POST-SURGERY: BREAST AUGMENTATION

INSTRUCTIONS

- Take your pain medication as directed.
- Ice packs will be applied in the recovery room. Continue to rotate them 30 minutes on, 30 minutes off while awake. Use them for the first 24-48 hours or longer for comfort.
- Begin your antibiotic the day after your surgery.
- Decreased activity and narcotic pain medication may cause constipation, increase fluid intake and use a stool softener as needed.
- Do not drink alcohol while taking pain medication.
- Sleep on your back with your head elevated for the first week.
- Substitute Tylenol (acetaminophen) for your narcotic pain medication as tolerated. For most this is within the first few days after surgery.
- Your compression bra should remain on 24 hours a day for the first week. A front closure sports bra may replace it for week 2-4. A regular bra of your choice can be worn after week 4.
- Your white compression stockings may be removed the morning after surgery.

ACTIVITIES

- Do not drive until you are no longer taking narcotic pain medication and have full range of motion with your arms.
- Start walking as soon as possible. Begin by walking around your house for the first few days and then increase activity slowly as your energy returns.
- Restrict excessive use of your arms for at least 5-7 days. Your elbows should not be lifted above your shoulders for the first week.
- To avoid excessive swelling or bleeding refrain from vigorous activities, anything that makes you feel excessively tired and no lifting anything heavier than 20 lbs for the first 2 weeks.
- Social and employment activities can be resumed in 5-7 days.
- Remember, if it hurts you are probably not ready to do it.

INCISION CARE

- Sutures are clear in appearance and will dissolve.
- Keep steri-strips covering your incisions clean and dry until they are removed in the office at one week post-surgery. Expect a small amount of discharge on the steri-strips.
- After steri-strips are removed you may shower normally and gently wash the incisions with your normal body wash or soap.
- Pat the incisions dry and expose to air for a few minutes each day.
- You may pad the incisions with gauze or a sanitary pad for comfort.
- Avoid exposing scars to sun for at least 12 months.
- The skin that has been stretched will burn easily so use a strong sun block



HEDIEH A. STEFANACCI, M.D.
AESTHETIC & RECONSTRUCTIVE PLASTIC SURGERY

INCISION CARE (continued)

- (SPF 30 or greater) if sun exposure is unavoidable.
- Do not wear an underwire bra until the incisions are completely healed, usually by 4 weeks.

WHAT TO EXPECT

- You may experience some numbness and/or a burning sensation of nipples and incision sites for about 2 weeks.
- It is normal to experience soreness, tightness, swelling and bruising for the first month after surgery. All patients heal at different rates so please be patient during this time.
- It is absolutely normal for your breasts to soften and settle at different rates so your final result may not be evident for several months.
- Your breasts may be sensitive to stimulation for a few weeks. This will resolve as you heal.
- Scars may be red and angry looking for 6 months or longer but usually soften and fade with time.

WHEN TO CALL

- If you have increased pain that is not relieved by medication.
- If you have any side effects to medications such as rash, nausea, vomiting or severe headache.
- If you have a temperature over 101 degrees.
- If you have bleeding from an incision that does not stop with light pressure.
- If you have yellowish or greenish drainage from the incisions or notice a foul odor.

FOR MEDICAL QUESTIONS, PLEASE CALL:

- (559) 797-9000, Monday-Thursday 9 a.m. – 5 p.m. and Friday 9 a.m.-noon.
- After hours and on weekends call the office number and Dr. Stefanacci will be paged.



HEDIEH A. STEFANACCI, M.D.
AESTHETIC & RECONSTRUCTIVE PLASTIC SURGERY